

Leading and higher twists in the proton polarized structure function g_1^p at large Bjorken- x ^a

Silvano Simula^(*), M. Osipenko^(**), G. Ricco^(***) and M. Taiuti^(***)

^(*) Istituto Nazionale di Fisica Nucleare, Sezione Roma III
Via della Vasca Navale 84, I-00146 Roma, Italy

^(**) Physics Department, Moscow State University, 119899 Moscow, Russia

^(***) University of Genova and Istituto Nazionale di Fisica Nucleare - Sezione di Genova
Via Dodecanneso 33, I-16146, Genova, Italy

Abstract

Power corrections to the Q^2 behavior of the Nachtmann moments of the proton polarized structure function g_1^p are investigated at large Bjorken x by developing a phenomenological fit of both the resonance (including the photon point) and deep inelastic data up to $Q^2 \sim 50 \text{ (GeV/c)}^2$. The leading twist is treated at NLO in the strong coupling constant and the effects of higher orders of the perturbative series are estimated using soft-gluon resummation techniques. In case of the first moment higher-twist effects are found to be quite small for $Q^2 \gtrsim 1 \text{ (GeV/c)}^2$, and the singlet axial charge is determined to be $a_0[10 \text{ (GeV/c)}^2] = 0.16 \pm 0.09$. In case of higher order moments, which are sensitive to the large- x region, higher-twist effects are significantly reduced by the introduction of soft gluon contributions, but they are still relevant at $Q^2 \sim \text{few (GeV/c)}^2$ at variance with the case of the unpolarized transverse structure function of the proton. This finding suggests that spin-dependent correlations among partons may have more impact than spin-independent ones. It is also shown that the parton-hadron local duality is violated in the region of polarized electroproduction of the $\Delta(1232)$ resonance.

^aProceedings of the IX International Conference on the *The Structure of Baryons*, Jefferson Lab (Newport News, USA), March 3-8, 2002, World Scientific Publishing (Singapore), in press.

1 Introduction

The experimental investigation of lepton deep-inelastic scattering (*DIS*) off proton and deuteron targets has provided a wealth of information on parton distributions in the nucleon. In the past few years some selected issues in the kinematical regions corresponding to large values of the Bjorken variable x have attracted a lot of theoretical and phenomenological interest; among them one should mention the occurrence of power corrections associated to *dynamical* higher-twist operators measuring the correlations among partons. The extraction of the latter is of particular relevance since the comparison with theoretical predictions either based on lattice *QCD* simulations or obtained from models of the nucleon structure represents an important test of *QCD* in its non-perturbative regime.

In Refs. [1] and [2] the world data on the unpolarized nucleon structure functions F_2^N and F_L^N have been used to carry out power correction analyses. In this contribution we summarize the main results [3] of the extension of such a twist analysis to the case of the polarized proton structure function g_1^p , performed in terms of Nachtmann moments. The latter however require the knowledge of the polarized structure function g_1^p in the whole x -range for fixed values of Q^2 , and therefore a new parameterization of g_1^p , which describes the *DIS* proton data up to $Q^2 \sim 50 \text{ (GeV/c)}^2$ and includes a phenomenological Breit-Wigner ansatz able to reproduce the existing electroproduction data in the proton-resonance regions, has been developed [3]. The interpolation formula for g_1^p has been successfully extended down to the photon point, showing that it nicely reproduces the very recent data [4] on the energy dependence of the asymmetry of the transverse photoproduction cross section as well as the experimental value of the proton Drell-Hearn-Gerasimov (*DHG*) sum rule. According to our parameterization of g_1^p the generalized *DHG* sum rule is predicted to have a zero-crossing point at $Q^2 = 0.16 \pm 0.04 \text{ (GeV/c)}^2$. Finally, the Q^2 behavior of low-order polarized Nachtmann moments has been obtained in the Q^2 -range between 0.5 and 50 (GeV/c)^2 .

2 Parton-hadron local duality in g_1^p

An important feature of the results obtained in Ref. [3] is that the resonant contribution to the polarized Nachtmann moments is negative for $Q^2 \sim \text{few (GeV/c)}^2$. This is mainly due to the well established fact [5] that the proton transverse asymmetry A_1^p in the $\Delta(1232)$ -resonance regions is negative up to $Q^2 \sim 3 \div 4 \text{ (GeV/c)}^2$. Thus, for $Q^2 \sim \text{few (GeV/c)}^2$ the resonant contribution to the polarized proton structure function is opposite in sign with respect to the unpolarized case (see Ref. [1]).

It is therefore legitimate to ask ourselves whether the parton-hadron local duality, observed empirically [6] in the unpolarized transverse structure function of the proton, holds as well in the polarized case. To this end we have generated pseudo-data in the resonance regions and in the *DIS* regime via our interpolation formula for g_1^p . Our results clearly shows that: i) at values of Q^2 as low as $\sim 0.5 \text{ (GeV/c)}^2$ there is no evidence at all of an occurrence of the local duality, as in the case of the unpolarized transverse structure function of the proton [7], and ii) in the kinematical regions where the $\Delta(1232)$ resonance is prominently

produced, the local duality breaks down at least for Q^2 up to few $(GeV/c)^2$, while in the higher resonance regions for $Q^2 \gtrsim 1 (GeV/c)^2$ it is not excluded by our parameterization. Note that in the unpolarized case the onset of the local duality occurs [6, 7] at $Q^2 \simeq 1 \div 2 (GeV/c)^2$, including also the $\Delta(1232)$ resonance regions. It should be mentioned that the usefulness of the concept of parton-hadron local duality relies mainly on the possibility to address the *DIS* curve at large x through measurements at low Q^2 in the resonance regions. It is therefore clear that the breakdown of the local duality in the region of the $\Delta(1232)$ resonance forbid us to get information from duality on the behavior of the scaling curve at the highest values of x .

3 Twist analysis of the polarized Nachtmann moments

In this Section we present the power correction analysis of the polarized Nachtmann moments, $M_n^{(1)}(Q^2)$, obtained in Ref. [3]. The leading twist, $\mu_n^{(1)}(Q^2)$, is treated both at next-to-leading (*NLO*) order and beyond any fixed order by adopting available soft gluon resummation (*SGR*) techniques. As for the power corrections, a phenomenological ansatz is considered, viz.

$$M_n^{(1)}(Q^2) = \mu_n^{(1)}(Q^2) + a_n^{(4)} \frac{\mu^2}{Q^2} \left[\frac{\alpha_s(Q^2)}{\alpha_s(\mu^2)} \right]^{\gamma_n^{(4)}} + a_n^{(6)} \left(\frac{\mu^2}{Q^2} \right)^2 \left[\frac{\alpha_s(Q^2)}{\alpha_s(\mu^2)} \right]^{\gamma_n^{(6)}} \quad (1)$$

where the logarithmic *pQCD* evolution of the twist-4 (twist-6) contribution is accounted for by an effective anomalous dimension $\gamma_n^{(4)}$ ($\gamma_n^{(6)}$) and the parameter $a_n^{(4)}$ ($a_n^{(6)}$) represents the overall strength of the twist-4 (twist-6) term at the renormalization scale μ^2 , chosen to be equal to $\mu^2 = 1 (GeV/c)^2$. In order to fix the running of the coupling constant $\alpha_s(Q^2)$, the updated *PDG* value $\alpha_s(M_Z^2) = 0.118$ is adopted.

In case of the first Nachtmann moment ($n = 1$) the corresponding leading twist term, $\mu_1^{(1)}(Q^2)$, does not receive any correction from *SGR* and therefore at *NLO* it reads as

$$\mu_1^{(1)}(Q^2) = \frac{\langle e^2 \rangle}{2} [\Delta q^{NS} + a_0(Q^2)] \left[1 - \frac{\alpha_s(Q^2)}{\pi} \right] \quad (2)$$

The non-singlet moment Δq^{NS} is taken fixed at the value $\Delta q^{NS} = 1.095$, deduced from the experimental values of the triplet and octet axial coupling constants, with the latter obtained under the assumption of *SU*(3)-flavor symmetry. The values of the singlet axial charge $a_0(\mu^2)$ and of the four higher-twist quantities $a_1^{(4)}$, $\gamma_1^{(4)}$, $a_1^{(6)}$ and $\gamma_1^{(6)}$ are determined by fitting our pseudo-data, adopting the least- χ^2 procedure in the Q^2 -range between 0.5 and 50 $(GeV/c)^2$. It turns out that the total contribution of the higher twists is tiny for $Q^2 \gtrsim 1 (GeV/c)^2$, but it is comparable with the leading twist already at $Q^2 \simeq 0.5 (GeV/c)^2$. Since the first moment basically corresponds to the area under the structure function g_1^p (as it is the case of the second moment of the unpolarized structure function F_2^p), the dominance of the leading twist in $M_1^{(1)}(Q^2)$, occurring for $Q^2 \gtrsim 1 (GeV/c)^2$, reflects only the concept of global duality

and *not* that of local duality (cf. Ref. [7]). In our analysis, where the leading and the higher twists are simultaneously extracted, the singlet axial charge (in the AB scheme) is determined to be $a_0(10 \text{ GeV}^2) = 0.16 \pm 0.09$, which nicely agrees with many recent estimates appeared in the literature. Our value of a_0 is therefore significantly below the naive quark-model expectation (i.e. compatible with the well known "proton spin crisis"), but it does not exclude completely a singlet axial charge as large as $\simeq 0.25$.

In case of higher-order moments ($n \geq 3$) both the NLO approximation and the SGR approach have been considered for the leading twist. The comparison of the corresponding twist analyses shows [3] that, except for the third moment, the contribution of the twist-2 is enhanced by soft gluon effects, while the total higher-twist term decreases significantly after the resummation of soft gluons. Thus, as already observed [2] in the unpolarized case, also in the polarized one it is mandatory to go beyond the NLO approximation and to include soft gluon effects in order to achieve a safer extraction of higher twists at large x , particularly for $Q^2 \sim \text{few } (\text{GeV}/c)^2$.

Finally, the twist decomposition of the polarized Nachtmann moments has been compared with the corresponding one of the unpolarized (transverse) Nachtmann moments obtained in Ref. [2] adopting the same SGR technique. It turns out [3] that the extracted higher-twist contribution appears to be a larger fraction of the leading twist in case of the polarized moments. This findings suggests that spin-dependent multiparton correlations may have more impact than spin-independent ones.

References

- [1] G. Ricco et al.: Nucl. Phys. **B555**, 306 (1999).
- [2] S. Simula: Phys. Lett. **B493**, 325 (2000).
- [3] S. Simula et al.: Phys. Rev. **D65**, 034017 (2002).
- [4] J. Ahrens et al.: Phys. Rev. Lett. **87**, 022003 (2001).
- [5] V. Burkert: Czech. J. Phys. **46**, 628 (1996).
- [6] E. Bloom and F. Gilman: Phys. Rev. Lett. **25**, 1140 (1970); Phys. Rev. **D4**, 2901 (1971).
- [7] G. Ricco et al.: Phys. Rev. **C57**, 356 (1998); Few-Body Syst. Suppl. **10**, 423 (1999). S. Simula: Phys. Lett. **B481**, 14 (2000); Phys. Rev. **D64**, 038301 (2001).